

LETTER TO THE EDITOR

Dear Editor

We came across a case of ectopic pregnancy in the adnexa that might be useful to share with our colleagues. A 38-year-old woman with history of infertility for 10 years was referred to our center. Upon admission, she had positive β HCG (with no prior infertility treatment), and expressed some pain in the left side of abdomen, but was generally in good conditions.



Fig 1. Transvaginal ultrasound of the septated uterus.




Fig 2. Transvaginal ultrasound of subserosal myoma.



Fig 3. Transvaginal ultrasound of ectopic pregnancy in the left adnex. One gestational sac is seen with two fetal poles and regular heart rates.

In transvaginal sonography, the uterus was septated with some subserosal myoma, and no gestational sac was evident in the uterine cavity. In the left adnex, however, one gestational sac with two fetal poles and regular fetal heart rates was seen (mono chorionic, monoamniotic). CRL measured 10mm, equivalent to 7 weeks' of gestation. The findings were compatible with ectopic pregnancy, and the patient underwent laparotomy and the embryos were completely removed. Figures 1–3 depict the transvaginal ultrasound images of the case.

F. Ahmadi MD¹

A. Vosough Taghi Dizaj MD¹

M. Salamati MD¹

Sh. Irani BSc²

1. Assistant Professor of Radiology, Department of Endocrinology and Female Infertility, Royan Institute, Tehran.

2. Bachelor of Science in Midwifery, Department of Endocrinology and Female Infertility, Royan Institute, Tehran.